

Please fax both pages of completed form to your team at 866.233.7151.

To reach your team, call toll-free 866.820.4844.

You can now monitor shipments and chat online if you have questions. Go to MyAccredoPatients.com to log in or get started.

Prescription & Enrollment Form Hereditary Angioedema (HAE)



Four simple steps to submit your referral.

Do not contact patient, benefits check only

1 Patient Information



Please provide copies of front and back of all medical and prescription insurance cards.

New patient Current patient

Patient's first name _____ Last name _____ Middle initial _____

Sex at birth: Male Female Preferred pronouns _____ Last 4 digits of SSN _____ Date of birth _____

Street address _____ Apt # _____

City _____ State _____ Zip _____

Home phone _____ Cell phone _____ Email address _____

Parent/guardian (if applicable) _____

Home phone _____ Cell phone _____ Email address _____

Alternate caregiver/contact _____

Home phone _____ Cell phone _____ Email address _____

OK to leave message with alternate caregiver/contact

Patient's primary language: English Other If other, please specify _____

Insurance Company _____

Phone _____ Identification # _____ Policy/group # _____

Prescription card: Yes No If yes, carrier _____ Policy #: _____ Group #: _____

Is patient eligible for Medicare? Yes No Does patient have secondary insurance? Yes No

2 Prescriber Information

All fields must be completed to expedite prescription fulfillment.

Date _____ Time _____ Date medication needed _____

Office/clinic/institution name _____

Prescriber info: Prescriber's first name _____ Last name _____

Prescriber's title _____ If NP or PA, under direction of Dr. _____

Office phone _____ Fax _____ NPI # _____ License # _____

Office contact and title _____ Office contact email _____

Office street address _____ Suite # _____

City _____ State _____ Zip _____

Infusion location: Patient's home Prescriber's office Infusion site If infusion site, complete information below dotted line:

Infusion info: Infusion site name _____ Clinic/hospital affiliation _____

Site street address _____ Suite # _____

City _____ State _____ Zip _____

Infusion site contact _____ Phone _____ Fax _____ Email _____

3 Clinical Information

Primary ICD-10 code (REQUIRED): _____ **D84.1 C1 esterase inhibitor [C1-INH] deficiency**

Other _____ Other drugs used to treat the disease _____

Weight _____ kg/lbs Date recorded _____ Height _____ cm/in Date recorded _____

NKDA Known drug allergies _____

Adverse reactions with previous HAE treatments? _____

If so, what brand of HAE caused the reaction? _____

Patient is naïve to HAE therapy Patient is continuing HAE therapy of _____

Concurrent meds _____

Patient's first name _____ Last name _____ Middle initial _____ Date of birth _____

Prescriber's first name _____ Last name _____ Phone _____

4 Prescribing Information

Medication	Strength/Formulation	Directions	Quantity/Refills
Cinryze (C1 Esterase Inhibitor [human])	500 unit vial	Infuse _____ units by slow IV injection at a rate of 1mL per minute every _____ days. Where clinically appropriate, please make dose divisible by 500 to avoid wastage.	Dispense: 1-month supply. Refill x 1 year unless noted otherwise Other _____
Berinerit (C1 Esterase Inhibitor [human])	500 unit vial	Infuse _____ units by slow IV injection at a rate of 4mL per minute as needed for acute hereditary angioedema (HAE) attack. Where clinically appropriate, please make dose divisible by 500 to avoid wastage.	Dispense: _____ doses. Keep at least _____ doses on hand at all times. Refill x 1 year unless noted otherwise. Other _____
Haegarda® (C1 Esterase Inhibitor Subcutaneous [human])—Fax mandatory hub form found here: https://accredo.com/prescribers/referral_forms/haegarda.pdf to 866.415.2162			
Ruconest (C1 Esterase Inhibitor [recombinant])—Fax mandatory hub form found here: https://accredo.com/prescribers/referral_forms/ruconest.pdf to 855.423.5757			
Takhzyro (lanadelumab-flyo)	300mg/2mL prefilled syringe	300mg by subcutaneous injection every two weeks 300mg by subcutaneous injection every four weeks	Dispense: 1-month supply. Refill x 1 year unless noted otherwise Other _____ Takhzyro should be administered by a healthcare provider or caregiver if that patient is under 12 years of age.
	150mg/1mL prefilled syringe	150mg by subcutaneous injection every two weeks 150mg by subcutaneous injection every four weeks	
icatibant	30mg/3mL prefilled syringe	Administer 30mg subcutaneously over at least 30 seconds for an acute attack of hereditary angioedema. If response is inadequate or symptoms recur, additional injections of 30mg may be administered at intervals of at least 6 hours. Do not administer more than 3 doses in 24 hours.	Dispense: _____ 30mg doses Keep at least three 30mg doses on hand at all times (unless noted otherwise _____ doses). Refill x 1 year unless noted otherwise Other _____
Kalbitor (ecallantide)	10mg/mL vial	Administer 30mg (3mL) subcutaneously in three 10mg (1mL) injections for an acute attack of hereditary angioedema. If the attack persists, may repeat the dose one time within a 24 hour period.	Dispense: Two 30mg doses. Keep at least two 30mg doses on hand at all times. Refill x 1 year unless noted otherwise Other _____

Kalbitor should only be administered by a healthcare professional with appropriate medical support to manage anaphylaxis.

Kalbitor to be infused in physician's office or controlled medical setting and/or Home infusion allowed by a Kalbitor trained RN

You must note the name of the brand product if brand is medically necessary for your patient _____

Infusion Requirements (for Cinryze, Berinerit and Kalbitor)

<p>Adverse reaction medications: (keep on hand at all times)</p> <p>Diphenhydramine 25mg by mouth or IV (for Kalbitor only) for mild allergic reactions and 50mg for moderate-severe.</p> <ul style="list-style-type: none"> <9kg: Diphenhydramine 1mg/kg up to max of 6.25mg 2–5 years old and >9kg: Diphenhydramine 6.25mg to 12.5mg 6–12 years old: Diphenhydramine 12.5mg to 25mg <p>Epinephrine 0.3mg auto-injector 2-pk for patient weighing greater than or equal to 30kg. Administer intramuscularly as needed for severe anaphylactic reaction times one dose; may repeat one time.</p> <p>Epinephrine 0.15mg auto-injector 2-pk, for patient weighing less than 30kg. Administer intramuscularly as needed for severe anaphylactic reaction times one dose; may repeat one time.</p> <p>For Kalbitor only: Normal Saline 250mL intravenously for anaphylactic reaction. Infuse wide open up to a max of 250mL per hour. Normal Saline 3mL flush before and after intravenous diphenhydramine administration and as needed for line patency.</p>	Refill x 1 year unless noted otherwise Other _____
---	---

Flushing orders (for Cinryze and Berinerit only):
 Normal saline 3mL intravenous (peripheral line) or 10mL intravenous (central line) before and after infusion, or as needed for line patency
 Heparin 100 units per mL 5mL intravenous (central line) as needed for final flush

Ancillary Supplies for all HAE products

Dispense needles, syringes and ancillary supplies necessary to administer medication.	Refill x 1 year unless noted otherwise Other _____
---	---

Nursing Start of Care Orders for all HAE products

Skilled nursing visit to provide patient education related to therapy, disease state, self and/or nurse administer of medication as prescribed. Visit frequency based on prescribed medication and dosage orders.

If shipped to physician's office or infusion clinic, physician accepts on behalf of patient for administration in office or infusion clinic.

Prescriber's signature required (sign below) (Physician attests this is his/her legal signature. NO STAMPS)

SIGN HERE Date _____ Dispense as written Date _____ Substitution allowed

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance with state-specific requirements could result in outreach to the prescriber.

The document(s) accompanying this transmission may contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender immediately and arrange for the return or destruction of these documents. All rights in the product names, trade names or logos of all third-party products that appear in this form, whether or not appearing with the trademark symbol, belong exclusively to their respective owners. © 2023 Accredo Health Group, Inc. I An Express Scripts Company. All rights reserved. HAE-00005-051023 CRP2405_8510

Prior Authorization Checklist Hereditary Angioedema (HAE)

Providing Accredo with the documentation outlined in this checklist may increase the likelihood and speed of obtaining coverage for your patients with HAE. Coverage criteria may vary by payer.

Referral Form	
	Completed HAE referral form ^{1,2}
	Copy of medical insurance card
	Copy of prescription benefits card
Clinical Documents	
	History of present illness (HAE)
	C1-inhibitor functional (or mutation) levels
	C4 antigenic levels
	Medication profile including any tried and failed therapies
Prescriber Specialization	
	Allergist
	Immunologist
	Hematologist
	Rheumatologist
	Other

Fax completed form to 866.233.7151.

If you have any questions, please call your Accredo provider support advocate, or call 866.820.4844.

¹For referral forms visit [accredo.com](https://www.accredo.com).

² Accredo referral form not required for electronic prescriptions or if using manufacturer hub form.