

Please fax both pages of completed form to your team at 888.302.1028.

To reach your team, call toll-free 844.516.3319.

You can now monitor shipments and chat online if you have questions. Go to MyAccredoPatients.com to log in or get started.

Prescription & Enrollment Form

Remicade[®] (infliximab) and Biosimilar

accredo[®]

Four simple steps to submit your referral.

1 Patient Information



Please provide copies of front and back of all medical and prescription insurance cards.

New patient Current patient

Patient's first name _____ Last name _____ Middle initial _____

Sex at birth: Male Female Preferred pronouns _____ Last 4 digits of SSN _____ Date of birth _____

Street address _____ Apt # _____

City _____ State _____ Zip _____

Home phone _____ Cell phone _____ E-mail address _____

Parent/guardian (if applicable) _____

Home phone _____ Cell phone _____ E-mail address _____

Alternate caregiver/contact _____

Home phone _____ Cell phone _____ E-mail address _____

OK to leave message with alternate caregiver/contact

Patient's primary language: English Other If other, please specify _____

2 Prescriber Information

All fields must be completed to expedite prescription fulfillment.

Date _____ Time _____ Date medication needed _____

Office/clinic/institution name _____

Prescriber info: Prescriber's first name _____ Last name _____

Prescriber's title _____ If NP or PA, under direction of Dr. _____

Office phone _____ Fax _____ NPI # _____ License # _____

Office contact and title _____ Office contact e-mail _____

Office street address _____ Suite # _____

City _____ State _____ Zip _____

Infusion location: Patient's home Prescriber's office Infusion site If infusion site, complete information below dotted line:

Infusion info: Infusion site name _____ Clinic/hospital affiliation _____

Site street address _____ Suite # _____

City _____ State _____ Zip _____

Infusion site contact _____ Phone _____ Fax _____ E-mail _____

3 Clinical Information

Primary ICD-10 code (REQUIRED): _____ Has the patient been treated previously for this condition? Yes No

Is patient currently on therapy? Yes No Please list all therapies tried/failed: _____

Patient wt _____ Date wt obtained _____

NKDA Known drug allergies _____

Concurrent meds _____

Patient's first name _____ Last name _____ Middle initial _____ Date of birth _____

Prescriber's first name _____ Last name _____ Phone _____

4 Prescribing Information

INFUSION LOCATION: Patient's home Healthcare facility

Medication	Directions	Quantity/Refills
Remicade® (infliximab) Inflectra® (infliximab-dyyb) Renflexis® (infliximab-abda) Avsola® (infliximab-axxq)	Loading dose: 5mg/kg _____ mg IV at week: 0, 2, 6 3mg/kg _____ mg IV at week: 0, 2, 6 Other _____ Maintenance dose: (_____ mg/kg) _____ mg IV every _____ weeks	Loading dose: 3 doses. No refills. Maintenance dose: 8-week supply. Refill x 1 year unless noted otherwise. _____ week supply Refill x 1 year unless noted otherwise. Other _____

Required medication and supplies for home infusion (please complete this section for home infusions only)

Premedication orders Acetaminophen 650mg PO 30 min prior to infusion; Diphenhydramine 50mg PO 30 min prior to infusion Other _____	Send quantity and refills sufficient for medication days supply
Infusion method: Infusion pump (If infusion pump checked, one will be provided) Gravity	
Fluids for administration and reconstitution (please strike through if not required) Fluid options should be as follows: NS 0.9% 250mL if dose 1000mg or less, NS 0.9% 500mL if dose > 1000mg Sterile Water as needed for reconstitution NS 0.9% Flush (if central venous access, sterile flush will be provided) Choose administration access: Peripheral access Central venous access If central venous access: Flush with 10mL Sterile NS 0.9% before and after infusion. Follow with heparin 100units/mL 5mL final flush If peripheral access: Flush with 3mL NS 0.9% before and after infusion and as needed	
Hypersensitivity/Anaphylaxis Stop infusion Medicate with: Epinephrine/EpiPen 0.3mg IM as needed for anaphylaxis (for children less than 30kg: Epinephrine 0.15mg) Start NS 0.9% 100mL at TKO Diphenhydramine 50mg slow IVP PRN anaphylaxis Hydrocortisone 100mg slow IVP PRN anaphylaxis Methylprednisolone 125mg slow IVP PRN anaphylaxis Diphenhydramine 50mg PO PRN anaphylaxis Other _____	
Skilled nursing visit as needed to establish venous access, administer medication and assess general status and response to therapy. *If nursing services will be required for therapy administration, the home health nurse will call for additional orders per state regulations.	
Lab orders _____ Frequency _____	

Dispense needles, syringes, ancillary supplies and home medical equipment necessary to administer medication.

If shipped to physician's office, physician accepts on behalf of patient for administration in office.

Prescriber's signature required (sign below) (Physician attests this is his/her legal signature. NO STAMPS)

SIGN HERE	_____	_____	_____
	Date	Dispense as written	Substitution allowed

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance with state-specific requirements could result in outreach to the prescriber.