

Please fax both pages of completed form to your team at 808.650.6487.

To reach your team, call toll-free 808.650.6488.

You can now monitor shipments and chat online if you have questions. Go to MyAccredoPatients.com to log in or get started.

Prescription & Enrollment Form
Dupixent® (dupilumab)

accredo®
677 Ala Moana Blvd., Suite 404,
Honolulu, HI 96813-5412

Four simple steps to submit your referral.

1 Patient Information



Please provide copies of front and back of all medical and prescription insurance cards.

New patient Current patient

Patient's first name _____ Last name _____ Middle initial _____

Preferred patient first name _____ Preferred patient last name _____

Sex at birth: Male Female Gender identity _____ Pronouns _____ Last 4 digits of SSN _____

Date of birth _____ Street address _____ Apt # _____

City _____ State _____ Zip _____

Home phone _____ Cell phone _____ Email address _____

Parent/guardian (if applicable) _____

Home phone _____ Cell phone _____ Email address _____

Alternate caregiver/contact _____

Home phone _____ Cell phone _____ Email address _____

OK to leave message with alternate caregiver/contact

Patient's primary language: English Other If other, please specify _____

2 Prescriber Information

All fields must be completed to expedite prescription fulfillment.

Date _____ Time _____ Date medication needed _____

Office/clinic/institution name _____

Prescriber info: Prescriber's first name _____ Last name _____

Prescriber's title _____ If NP or PA, under direction of Dr. _____

Office phone _____ Fax _____ NPI # _____ License # _____

Office contact and title _____ Office contact email _____

Office street address _____ Suite # _____

City _____ State _____ Zip _____

Infusion location: Patient's home Prescriber's office Infusion site If infusion site, complete information below dotted line:

Infusion info: Infusion site name _____ Clinic/hospital affiliation _____

Site street address _____ Suite # _____

City _____ State _____ Zip _____

Infusion site contact _____ Phone _____ Fax _____ Email _____

3 Clinical Information

ICD-10 code (REQUIRED): _____

NKDA Known drug allergies _____

Prior anaphylactic reaction: Yes (Reason/date _____) No

Concurrent meds _____ Estimated % BSA involvement _____

Concomitant therapies: Short-acting beta agonist Long-acting beta agonist Antihistamines Decongestants Immunotherapy

Inhaled corticosteroid Leukotriene modifiers Oral steroids Nasal steroids Other _____

Lab results: History of positive skin OR RAST test to a perennial aeroallergen

Pre-treatment steroid dose _____ mg Pre-treatment serum IgE level _____ IU per mL Test date _____

Pre-treatment serum eosinophils _____ cells/mcL and/or sputum eosinophils _____ Date _____

Patient wt _____ kg Date wt obtained _____

MD Specialty (required): Allergist Pulmonologist ENT Primary care Pediatrician Dermatologist Other _____

Prescription type: Naïve/new start Restart Continued therapy

Prior therapies: Please fax detailed medication history with dates of use as available. Required by some plan authorization criteria.

Topical steroid(s) Oral antihistamines Topical PDE-4 inhibitor Oral steroids Oral immunosuppressants

Topical calcineurin inhibitor Sinus surgery

Patient's first name _____ Last name _____ Middle initial _____ Date of birth _____

Prescriber's first name _____ Last name _____ Phone _____

4 Prescribing Information

Medication	Strength / Formulation and Directions	Quantity/Refills	
Dupixent® (dupilumab) 200mg/1.14mL pre-filled pen 2-pack Dupixent® (dupilumab) 200mg/1.14mL pre-filled syringe 2-pack Dupixent® (dupilumab) 300mg/2mL pre-filled pen 2-pack Dupixent® (dupilumab) 300mg/2mL pre-filled syringe 2-pack	Starter Dose: Inject 400mg under the skin on Day 1 then 200mg every 2 weeks starting on day 15 and thereafter. Maintenance Dose: Inject 200mg under the skin every 2 weeks.	Starter dose: Quantity _____ No refills	
	Starter Dose: Inject 600mg under the skin on Day 1 then 300mg every 2 weeks starting on day 15 and thereafter. Maintenance Dose: Inject 300mg under the skin every 2 weeks.	Maintenance dose: Quantity _____ Refills _____	
	Starter Dose: Inject 600mg under the skin on Day 1 then 300mg every 4 weeks thereafter starting on day 29. Maintenance Dose: Inject 300mg under the skin every 4 weeks.	For indications without a starter dose: Inject 100mg under the skin every 2 weeks Inject 200mg under the skin every 2 weeks Inject 200mg under the skin every 4 weeks Inject 300mg under the skin once weekly Inject 300mg under the skin every 2 weeks Inject 300mg under the skin every 4 weeks	For indications without a starter dose: Quantity _____ Refills _____
			Patient weight _____ kg

If shipped to physician's office or infusion clinic, physician accepts on behalf of patient for administration in office or infusion clinic.

Prescriber's signature required (sign below) (Physician attests this is his/her legal signature. NO STAMPS)

**SIGN
HERE**

_____ **Date**

_____ **Dispense as written**

_____ **Date**

_____ **Substitution allowed**

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance with state-specific requirements could result in outreach to the prescriber.