

Please fax both pages of completed form to the Psoriasis team at 808.650.6487.

To reach your team, call toll-free 808.650.6488.

You can now monitor shipments and chat online if you have questions. Go to [MyAccredoPatients.com](http://MyAccredoPatients.com) to log in or get started.

## Prescription & Enrollment Form Psoriasis

*accredo*<sup>®</sup>  
677 Ala Moana Blvd., Suite 404,  
Honolulu, HI 96813-5412

Four simple steps to submit your referral.

### 1 Patient Information



Please provide copies of front and back of all medical and prescription insurance cards.

New patient    Current patient

Patient's first name \_\_\_\_\_ Last name \_\_\_\_\_ Middle initial \_\_\_\_\_

Preferred patient first name \_\_\_\_\_ Preferred patient last name \_\_\_\_\_

Sex at birth:    Male    Female    Gender identity \_\_\_\_\_    Pronouns \_\_\_\_\_    Last 4 digits of SSN \_\_\_\_\_

Date of birth \_\_\_\_\_ Street address \_\_\_\_\_ Apt # \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Home phone \_\_\_\_\_ Cell phone \_\_\_\_\_ Email address \_\_\_\_\_

Parent/guardian (if applicable) \_\_\_\_\_

Home phone \_\_\_\_\_ Cell phone \_\_\_\_\_ Email address \_\_\_\_\_

Alternate caregiver/contact \_\_\_\_\_

Home phone \_\_\_\_\_ Cell phone \_\_\_\_\_ Email address \_\_\_\_\_

OK to leave message with alternate caregiver/contact

Patient's primary language:    English    Other    If other, please specify \_\_\_\_\_

### 2 Prescriber Information

All fields must be completed to expedite prescription fulfillment.

Date \_\_\_\_\_ Time \_\_\_\_\_ Date medication needed \_\_\_\_\_

Office/clinic/institution name \_\_\_\_\_

**Prescriber info:** Prescriber's first name \_\_\_\_\_ Last name \_\_\_\_\_

Prescriber's title \_\_\_\_\_ If NP or PA, under direction of Dr. \_\_\_\_\_

Office phone \_\_\_\_\_ Fax \_\_\_\_\_ NPI # \_\_\_\_\_ License # \_\_\_\_\_

Office contact and title \_\_\_\_\_ Office contact email \_\_\_\_\_

Office street address \_\_\_\_\_ Suite # \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Infusion location:    Patient's home    Prescriber's office    Infusion site    If infusion site, complete information below dotted line:  
-----

**Infusion info:** Infusion site name \_\_\_\_\_ Clinic/hospital affiliation \_\_\_\_\_

Site street address \_\_\_\_\_ Suite # \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Infusion site contact \_\_\_\_\_ Phone \_\_\_\_\_ Fax \_\_\_\_\_ Email \_\_\_\_\_

### 3 Clinical Information

**Primary ICD-10 code (REQUIRED):** \_\_\_\_\_ Severity:    Moderate    Moderate to severe    Severe    BSA \_\_\_\_\_%

Type:    Plaque    Other \_\_\_\_\_

Significant symptoms \_\_\_\_\_

Prior Treatments:    Topicals    PUVA    UVB    Methotrexate    Cyclosporine    Oral retinoid    Other \_\_\_\_\_

Medical justification for prescribing \_\_\_\_\_

NKDA    Known drug allergies \_\_\_\_\_

Concurrent meds \_\_\_\_\_

Patient's first name \_\_\_\_\_ Last name \_\_\_\_\_ Middle initial \_\_\_\_\_ Date of birth \_\_\_\_\_

Prescriber's first name \_\_\_\_\_ Last name \_\_\_\_\_ Phone \_\_\_\_\_

## 4 Prescribing Information

Medication	Strength/Formulation	Directions	Quantity/Refills
Bimzelx® (bimekizumab-bkzx)	160mg Prefilled Syringe (PFS) 160mg Autoinjector	<b>Loading dose:</b> 320mg (given as two 160mg injections) at Weeks 0, 4, 8, 12, and 16, then every 8 weeks thereafter.	QS for 1-month 4 Refills
		<b>Maintenance dose:</b> Inject 320mg subcutaneously every 8 weeks. Inject 320mg subcutaneously every 4 weeks (for patients weighing ≥ 120kg)	QS for 1 dose Other _____ Refills _____
Cimzia® (certolizumab)	200mg/mL PFS 200mg/mL Lyophilized Powder in Single Dose vial for Reconstitution	<b>Loading dose:</b> Inject 400mg subcutaneously at weeks 0, 2 and 4.	1 starter kit OR- QS for 1-month loading dose No Refills
		<b>Maintenance dose:</b> Inject 400mg subcutaneously every 2 weeks. Inject 200mg subcutaneously every 2 weeks. Other _____	1-month supply 3-month supply Refill QS 1 year unless otherwise noted Other _____
Cosentyx® (secukinumab)	75mg PFS 150mg PFS 150mg Pen 300mg (2x150mg) PFS 300mg (2x150mg) Pen 300mg Unoready Pen	<b>Loading dose:</b> Inject _____mg subcutaneously at weeks 0, 1, 2, 3 and 4 followed by _____ every 4 weeks.	QS for 5 doses No Refills
		<b>Maintenance dose:</b> Inject _____mg subcutaneously every 4 weeks.	1-month supply 3-month supply Refill QS 1 year unless otherwise noted Other _____
Enbrel® (etanercept)	25mg Single Use vial 25mg PFS 50mg PFS 50mg SureClick™ 50mg Mini Cartridge	<b>Loading dose:</b> Inject 50mg subcutaneously twice a week x 3 months, then 50mg once a week.	QS for 3-month loading dose No Refills
		<b>Maintenance dose:</b> Inject 50mg subcutaneously once a week. Inject _____mg subcutaneously _____ per week.	1-month supply 3-month supply Refill QS 1 year unless otherwise noted Other _____
Other _____			
<b>Ancillary Supplies: (Prescriber to strike through if not required)</b> Dispense ancillary supplies such as needles, syringes, sterile water, etc. and home medical equipment necessary to administer the therapy as needed.			Send quantity sufficient for medication days supply

If shipped to physician's office or infusion clinic, physician accepts on behalf of patient for administration in office or infusion clinic.  
By signing below, I certify that the above therapy is medically necessary.

Prescriber's signature required (sign below) (Physician attests this is his/her legal signature. NO STAMPS)

**SIGN  
HERE**

Date

Dispense as written

Date

Substitution allowed

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance with state-specific requirements could result in outreach to the prescriber.

Patient's first name \_\_\_\_\_ Last name \_\_\_\_\_ Middle initial \_\_\_\_\_ Date of birth \_\_\_\_\_

Prescriber's first name \_\_\_\_\_ Last name \_\_\_\_\_ Phone \_\_\_\_\_

## 4 Prescribing Information

Medication	Strength/Formulation	Directions	Quantity/Refills
Ilumya™ (tildrakizumab-asmn)	100mg/mL in a single-dose PFS	<b>Loading dose:</b> Inject 100mg subcutaneously at weeks 0, 4 and every 12 weeks thereafter.	2 syringes for loading/ induction dose No Refills
		<b>Maintenance dose:</b> Inject 100mg subcutaneously every 12 weeks.	1-month supply 3-month supply Refill QS 1 year unless otherwise noted Other _____
Otezla® (apremilast)	Starter Pack (28 day)	<b>Loading dose:</b> Day 1: Take by mouth 10mg in the morning. Day 2: Take by mouth 10mg in the morning and 10mg in the evening. Day 3: Take by mouth 10mg in the morning and 20mg in the evening. Day 4: Take by mouth 20mg in the morning and 20mg in the evening. Day 5: Take by mouth 20mg in the morning and 30mg in the evening. Day 6 and thereafter: Take by mouth 30mg in the morning and 30mg in the evening.	1 Kit No Refills
	30mg tablets	<b>Maintenance dose:</b> Take 30mg by mouth twice a day. Take 30mg by mouth once a day (severe renal impairment).	1-month supply 3-month supply Refill QS 1 year unless otherwise noted Other _____
Siliq™ (brodalumab)	210mg/1.5mL prefilled syringe (PFS) (2-pack)	<b>Loading dose:</b> Inject 210mg subcutaneously at weeks 0, 1 and 2 followed by 210mg every 2 weeks.	2 Kits No Refills
		<b>Maintenance dose:</b> Inject 210mg subcutaneously every 2 weeks.	1-month supply 3-month supply Refill QS 1 year unless otherwise noted Other _____
Skyrizi™ (risankizumab-rzaa)	150mg/mL in each single-dose PFS 150mg/mL in each single-dose pen	<b>Loading dose:</b> Inject 150mg subcutaneously at weeks 0, 4 and every 12 weeks thereafter.	2 doses for loading/ induction No Refills
		<b>Maintenance dose:</b> Inject 150mg subcutaneously every 12 weeks.	1-month supply 3-month supply Refill QS 1 year unless otherwise noted Other _____
Other _____			
<b>Ancillary Supplies: (Prescriber to strike through if not required)</b> Dispense ancillary supplies such as needles, syringes, sterile water, etc. and home medical equipment necessary to administer the therapy as needed.			Send quantity sufficient for medication days supply

If shipped to physician's office or infusion clinic, physician accepts on behalf of patient for administration in office or infusion clinic.

**Prescriber's signature required (sign below) (Physician attests this is his/her legal signature. NO STAMPS)**

**SIGN  
HERE**

\_\_\_\_\_ Date

\_\_\_\_\_ Dispense as written

\_\_\_\_\_ Date

\_\_\_\_\_ Substitution allowed

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance with state-specific requirements could result in outreach to the prescriber.

Patient's first name \_\_\_\_\_ Last name \_\_\_\_\_ Middle initial \_\_\_\_\_ Date of birth \_\_\_\_\_

Prescriber's first name \_\_\_\_\_ Last name \_\_\_\_\_ Phone \_\_\_\_\_

## 4 Prescribing Information

Medication	Strength/Formulation	Directions	Quantity/Refills
Sotyktu™ (deucravacitinib)	6mg tablet	Take 1 tablet daily	Refill QS 1 year unless noted otherwise Other _____
Stelara® (ustekinumab)	45mg/0.5mL single-dose vial 45mg/0.5mL PFS 90mg/1mL PFS  Please include patient weight: _____ kg	<b>Loading dose:</b> Inject _____ mg subcutaneously at week 0 and week 4, followed by every 12 weeks thereafter	2 doses for loading/induction No Refills
		<b>Maintenance dose:</b> Inject _____ mg subcutaneously every 12 weeks	1-month supply 3-month supply Refill QS 1 year unless otherwise noted Other _____
Taltz® (ixekizumab)	80mg single-dose autoinjector 80mg single-dose PFS	<b>Loading and Induction dose:</b> Inject 160mg (two 80mg injections) subcutaneously at week 0, followed by 80mg at weeks 2, 4, 6, 8, 10 and 12, then 80mg every 4 weeks.	8 devices for loading/induction No Refills
		<b>Maintenance dose:</b> Inject 80mg subcutaneously every 4 weeks	1-month supply 3-month supply Refill QS 1 year unless otherwise noted Other _____
Tremfya™ (guselkumab)	100mg/mL in each single-dose PFS 100mg/mL in each single-dose pen	<b>Loading dose:</b> Inject 100mg subcutaneously at weeks 0, 4 and every 8 weeks thereafter.	2 doses for loading/induction No Refills
		<b>Maintenance dose:</b> Inject 100mg subcutaneously every 8 weeks.	1-month supply 3-month supply Refill QS 1 year unless otherwise noted Other _____
Other _____			
<b>Ancillary Supplies: (Prescriber to strike through if not required)</b> Dispense ancillary supplies such as needles, syringes, sterile water, etc. and home medical equipment necessary to administer the therapy as needed.			Send quantity sufficient for medication days supply

If shipped to physician's office or infusion clinic, physician accepts on behalf of patient for administration in office or infusion clinic.

**Prescriber's signature required (sign below) (Physician attests this is his/her legal signature. NO STAMPS)**

**SIGN  
HERE**

Date \_\_\_\_\_

Dispense as written

Date \_\_\_\_\_

Substitution allowed

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance with state-specific requirements could result in outreach to the prescriber.