

Please fax both pages of completed form to your team at 808.650.6487.

To reach your team, call toll-free 808.650.6488.

You can now monitor shipments and chat online if you have questions. Go to MyAccredoPatients.com to log in or get started.

Prescription & Enrollment Form

Cystic fibrosis—oral

accredo®

677 Ala Moana Blvd., Suite 404,
Honolulu, HI 96813-5412

Four simple steps to submit your referral.

1 Patient Information



Please provide copies of front and back of all medical and prescription insurance cards.

New patient Current patient

Patient's first name _____ Last name _____ Middle initial _____

Sex at birth: Male Female Pronouns _____ Last 4 digits of SSN _____ Date of birth _____

Street address _____ Apt # _____

City _____ State _____ Zip _____

Home phone _____ Cell phone _____ Email address _____

Parent/guardian (if applicable) _____

Home phone _____ Cell phone _____ Email address _____

Alternate caregiver/contact _____

Home phone _____ Cell phone _____ Email address _____

OK to leave message with alternate caregiver/contact

Patient's primary language: English Other If other, please specify _____

2 Prescriber Information

All fields must be completed to expedite prescription fulfillment.

Date _____ Time _____ Date medication needed _____

Office/clinic/institution name _____

Prescriber's first name _____ Last name _____

Prescriber's title _____ If NP or PA, under direction of Dr. _____

Office phone _____ Fax _____ NPI # _____ License # _____

Office contact and title _____ Office contact email _____

Office street address _____ Suite # _____

City _____ State _____ Zip _____

Deliver product to: Prescriber's office Patient's home

3 Clinical Information

Primary ICD-10 code (REQUIRED): _____

Weight _____ kg/lbs Height _____ cm/in Date recorded _____

CFR Mutation type(s): F508del G551D G1244E G1349D G178R G551S S1251N
 S1255P S549N S549R R117H Other _____

Patient is: Heterozygous Homozygous for above mutation(s) FEV 1 _____ Date _____

NKDA Known drug allergies _____

Concurrent meds _____

Baseline eye exam date _____ Last hearing screen _____

Serum Creatinine _____ Date _____ Estimated GFR _____

Patient's first name _____ Last name _____ Middle initial _____ Date of birth _____

Prescriber's first name _____ Last name _____ Phone _____

4 Prescribing Information

Medication	Strength/Formulation	Directions	Quantity/Refills
Mutation Correctors			
Kalydeco® (ivacaftor) tablets	(ages 6 years and older) 150mg tablet	Take 1 tablet by mouth every 12 hours with fat-containing food. Other _____ <i>(i.e. dose adjustments for hepatic impairment and moderate to strong CYP3A inhibitors; please see package insert.)</i>	1-month supply 3-month supply Other _____
Kalydeco® (ivacaftor) oral granules	(ages 1 month–5 years) 5.8mg packet (aged 1–2 months; > 3kg) 13.4mg packet (aged 2–4 months; > 3kg) 25mg packet (aged 4–6 months; > 5kg) 25mg packet (aged > 6 months–5 years; 5-7kg) 50mg packet (aged > 6 months–5 years; 7-14kg) 75mg packet (aged > 6 months–5 years; > 14kg) Patient weight _____	Mix 1 packet of granules in one teaspoon (5mL) of soft food or liquid and administer every 12 hours with fat-containing food. Other _____ <i>(i.e. dose adjustments for hepatic impairment and moderate to strong CYP3A inhibitors; please see package insert.)</i>	Refills _____
Orkambi® (lumacaftor/ ivacaftor) tablet	(ages 6–11 years) 100mg/125mg tablet (12 years and older) 200mg/125mg tablet	Take 2 tablets by mouth every 12 hours with fat-containing food. Other _____ <i>(i.e. dose adjustments for hepatic impairment and moderate to strong CYP3A inhibitors; please see package insert.)</i>	
Orkambi® (lumacaftor/ ivacaftor) oral granules	(ages 1–5 years) 75mg/94mg granules (weight 7–9kg) 100mg/125mg granules (weight 9–14kg) 150mg/188mg granules (weight >14kg) Patient weight _____	Mix 1 packet of granules in one teaspoon (5mL) of soft food or liquid and administer every 12 hours with fat-containing food. Other _____ <i>(i.e. dose adjustments for hepatic impairment and moderate to strong CYP3A inhibitors; please see package insert.)</i>	
Symdeko® (tezacaftor/ ivacaftor + ivacaftor) tablets	50mg/75mg tablet + 75mg tablet 100mg/150mg tablet + 150mg tablet	Take 1 white tablet in the morning, and 1 blue tablet in the evening approximately 12 hours apart with fat-containing food. Take 1 yellow tablet by mouth in the morning, and 1 blue tablet in the evening approximately 12 hours apart with fat-containing food. Other _____ <i>(i.e. dose adjustments for hepatic impairment and moderate to strong CYP3A inhibitors; please see package insert.)</i>	
Trikafta® (elexacaftor/ tezacaftor/ ivacaftor + ivacaftor) tablets	(ages 6 years and older) 50mg/25mg/37.5mg tablet + 75mg tablet 100mg/50mg/75mg tablet + 150mg tablet Patient weight _____	Take 2 orange tablets by mouth in the morning, and 1 blue tablet in the evening approximately 12 hours apart with fat-containing food. Other _____ <i>(i.e. dose adjustments for hepatic impairment and moderate to strong CYP3A inhibitors; please see package insert.)</i>	
Trikafta® (elexacaftor/ tezacaftor/ ivacaftor+ ivacaftor) granules	(ages 2–5 years) 80mg/40mg/60mg +59.5mg oral granules (weight < 14kg) 100mg/50mg/75mg +75mg oral granules (weight > 14kg) Patient weight _____	Mix 1 blue packet in one teaspoon (5mL) of soft food or liquid and take in the morning. Mix 1 green packet in one teaspoon (5mL) of soft food or liquid and take in the evening. Take with fat-containing food approximately 12 hours apart. Mix 1 orange packet in one teaspoon (5mL) of soft food or liquid and take in the morning. Mix 1 pink packet in one teaspoon (5mL) of soft food or liquid and take in the evening. Take with fat-containing food approximately 12 hours apart. Other _____ <i>(i.e. dose adjustments for hepatic impairment and moderate to strong CYP3A inhibitors; please see package insert.)</i>	
Other			

If shipped to physician's office, physician accepts on behalf of patient for administration in office.

Prescriber's signature required (sign below) (Physician attests this is his/her legal signature. NO STAMPS)

**SIGN
HERE**

Date

Dispense as written

Date

Substitution allowed

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance with state-specific requirements could result in outreach to the prescriber.