

Please fax both pages of completed form to your team at 888.302.1028.

To reach your team, call toll-free 844.516.3319.

You can now monitor shipments and chat online if you have questions. Go to MyAccredoPatients.com to log in or get started.

Prescription & Enrollment Form

Arthritis and Inflammatory—Intravenous

accredo®

Four simple steps to submit your referral.

1 Patient Information



Please provide copies of front and back of all medical and prescription insurance cards.

New patient Current patient

Patient's first name _____ Last name _____ Middle initial _____

Sex at birth: Male Female Preferred pronouns _____ Last 4 digits of SSN _____ Date of birth _____

Street address _____ Apt # _____

City _____ State _____ Zip _____

Home phone _____ Cell phone _____ E-mail address _____

Parent/guardian (if applicable) _____

Home phone _____ Cell phone _____ E-mail address _____

Alternate caregiver/contact _____

Home phone _____ Cell phone _____ E-mail address _____

OK to leave message with alternate caregiver/contact

Patient's primary language: English Other If other, please specify _____

2 Prescriber Information

All fields must be completed to expedite prescription fulfillment.

Date _____ Time _____ Date medication needed _____

Office/clinic/institution name _____

Prescriber info: Prescriber's first name _____ Last name _____

Prescriber's title _____ If NP or PA, under direction of Dr. _____

Office phone _____ Fax _____ NPI # _____ License # _____

Office contact and title _____ Office contact e-mail _____

Office street address _____ Suite # _____

City _____ State _____ Zip _____

Infusion location: Patient's home Prescriber's office Infusion site If infusion site, complete information below dotted line:

Infusion info: Infusion site name _____ Clinic/hospital affiliation _____

Site street address _____ Suite # _____

City _____ State _____ Zip _____

Infusion site contact _____ Phone _____ Fax _____ E-mail _____

3 Clinical Information

Primary ICD-10 code (REQUIRED): _____

Has the patient been treated previously for this condition? Yes No Is patient currently on therapy? Yes No

Please list all therapies tried/failed: _____

Patient wt _____ Date wt obtained _____

NKDA Known drug allergies _____

Concurrent meds _____

Patient's first name _____ Last name _____ Middle initial _____ Date of birth _____

Prescriber's first name _____ Last name _____ Phone _____

4 Prescribing Information

Medication	Dose/Directions	Fluids for administration and reconstitution (please strike through if not required)	Quantity/Refills
Actemra® (tocilizumab)	<p>Rheumatoid Arthritis (RA): 4mg/kg intravenous infusion every 4 weeks. Maximum dose of 800mg/infusion 8mg/kg intravenous infusion every 4 weeks. Maximum dose of 800mg/infusion</p> <p>Polyarticular Juvenile Idiopathic Arthritis (PJIA): 10mg/kg intravenous infusion every 4 weeks (2 years or older, Less than 30kg) 8mg/kg intravenous infusion every 4 weeks (2 years or older, 30kg or greater)</p> <p>Systemic Juvenile Idiopathic Arthritis (SJIA) and Cytokine Release Syndrome: 12mg/kg intravenous infusion every 2 weeks (2 years or older, Less than 30kg) Maximum dose of 800mg/infusion 8mg/kg intravenous infusion every 2 weeks (2 years or older, 30kg or greater) Maximum dose of 800mg/infusion</p>	<p>Dilute desired dose with normal saline to total desired volume to be infused over 1 hour.</p> <p>NS 0.9% 100mL >30kg NS 0.9% 50mL < 30kg</p>	<p>Dispense 1-month supply. Refill x 1 year unless noted otherwise.</p> <p>Dispense 90-day supply. Refill x 1 year unless noted otherwise.</p> <p>Other _____</p> <p>Refills _____</p>
Orencia® (abatacept)	<p>Rheumatoid Arthritis and Psoriatic Arthritis: 500mg (less than 60kg) intravenous infusion 750mg (60 to 100kg) intravenous infusion 1000mg (over 100kg) intravenous infusion</p> <p>Juvenile Idiopathic Arthritis: 10mg/kg intravenous infusion (if less than 75kg) 750mg intravenous infusion (75 to 100kg) 1,000mg intravenous infusion (over 100kg) Starting dose: at week: 0, 2 and 4, then every 4 weeks</p>	<p>Reconstitute each vial of Orencia with 10mL of sterile water. Dilute desired dose to total of 100mL in normal saline to be infused over 30 minutes.</p> <p>NS 0.9% 100mL Sterile Water as needed for reconstitution.</p>	<p>Starter dose: x 3 doses. No refills.</p>
	<p>Maintenance dose: every 4 weeks</p>		<p>Maintenance dose: 1-month supply Refill x 1 year unless noted otherwise Other _____ Refills _____</p>
Simponi Aria® (golimumab)	<p>Starting dose: 2mg/kg _____ mg IV at week: 0, 4 and every 8 weeks Other _____</p>	<p>Dilute desired dose with normal saline to a total volume of 100mL to be infused over 30 minutes.</p>	<p>Starter dose: x 3 doses. No refills.</p>
	<p>Maintenance dose: 2mg/kg _____ mg IV every 8 weeks Other _____</p>		<p>Maintenance dose: 1-month supply Refill x 1 year unless noted otherwise Other _____ Refills _____</p>
Other			<p>Other _____ Refills _____</p>

If shipped to physician's office or infusion clinic, physician accepts on behalf of patient for administration in office or infusion clinic.

Prescriber's signature required (sign below) (Physician attests this is his/her legal signature. NO STAMPS)

**SIGN
HERE**

Date

Dispense as written

Date

Substitution allowed

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance with state-specific requirements could result in outreach to the prescriber.

