

Please fax all pages of completed form to your team at 888.302.1028.

To reach your team, call toll-free 844.516.3319.

You can now monitor shipments and chat online if you have questions. Go to MyAccredoPatients.com to log in or get started.

Accredo® Specialty Pharmacy Prescription & Enrollment Form

Tremfya®

EVERNORTH®
HEALTH SERVICES

Four simple steps to submit your referral.

1 Patient Information



Please provide copies of front and back of all medical and prescription insurance cards.

New patient Current patient

Patient's first name _____ Last name _____ Middle initial _____

Preferred patient first name _____ Preferred patient last name _____

Sex at birth: Male Female Gender identity _____ Pronouns _____ Last 4 digits of SSN _____

Date of birth _____ Street address _____ Apt # _____

City _____ State _____ Zip _____

Home phone _____ Cell phone _____ Email address _____

Parent/guardian (if applicable) _____

Home phone _____ Cell phone _____ Email address _____

Alternate caregiver/contact _____

Home phone _____ Cell phone _____ Email address _____

OK to leave message with alternate caregiver/contact

Patient's primary language: English Other If other, please specify _____

Provider will read the following statement to patient: By providing the phone number(s) and email address above, you consent to receiving automated/artificial voice calls, emails and/or text messages from Accredo about your prescription(s), account, and health care. Standard data rates apply. Message frequency varies.

2 Prescriber Information

All fields must be completed to expedite prescription fulfillment.

Date _____ Time _____ Date medication needed _____

Office/clinic/institution name _____

Prescriber info: Prescriber's first name _____ Last name _____

Prescriber's title _____ If NP or PA, under direction of Dr. _____

Office phone _____ Fax _____ NPI # _____ License # _____

Office contact and title _____ Office contact email _____

Office street address _____ Suite # _____

City _____ State _____ Zip _____

Infusion location: Patient's home Prescriber's office Infusion site If infusion site, complete information below dotted line: _____

Infusion info: Infusion site name _____ Clinic/hospital affiliation _____

Site street address _____ Suite # _____

City _____ State _____ Zip _____

Infusion site contact _____ Phone _____ Fax _____ Email _____

3 Clinical Information

Primary ICD-10 code (REQUIRED): _____ Has the patient been treated previously for this condition? Yes No

Is patient currently on therapy? Yes No Please list all therapies tried/failed: _____

Patient wt _____ Date wt obtained _____

NKDA Known drug allergies _____

Concurrent meds _____

Patient's first name _____ Last name _____ Middle initial _____ Date of birth _____

Prescriber's first name _____ Last name _____ Phone _____

4 Prescribing Information

Medication	Strength/Formulation	Directions	Quantity/Refills
Tremfya®	Psoriasis and Psoriatic Arthritis		
	100mg/mL in each single-dose One-Press injector 100mg/mL in each single-dose prefilled syringe (PFS)	Loading Dose: Inject 100mg subcutaneously at weeks 0, 4 and every 8 weeks thereafter	QS for loading period No Refills
		Maintenance Dose: Inject 100mg subcutaneously every 8 weeks	1-month supply 3-month supply Other _____ Refills _____
	Ulcerative Colitis		
	Loading: 200mg/20mL single-dose vial 200mg/2mL starter pack pen	Loading Dose: Infuse 200mg IV at weeks 0, 4 and 8 Inject 400mg (2-200mg/2mL) subcutaneously at weeks 0, 4 and 8	QS for loading period No Refills
	Maintenance: 100mg/mL in each single-dose One-Press injector 100mg/mL in each single-dose PFS 100mg/mL in each single-dose pen 200mg/2mL in each single-dose PFS 200mg/2mL in each single-dose pen	Maintenance Dose: Inject 100mg subcutaneously at week 16 and every 8 weeks thereafter Inject 200mg subcutaneously at week 12 and every 4 weeks thereafter Inject 100mg subcutaneously every 8 weeks Inject 200mg subcutaneously every 4 weeks	1-month supply 3-month supply Other _____ Refills _____
Other	Crohn's		
	Loading: 200mg/20mL single-dose vial 200mg/2mL starter pack pen	Loading Dose: Infuse 200mg IV at weeks 0, 4 and 8 Inject 400mg (2-200mg/2mL) subcutaneously at weeks 0, 4 and 8	QS for loading period No Refills
	Maintenance: 100mg/mL in each single-dose One-Press injector 100mg/mL in each single-dose PFS 100mg/mL in each single-dose pen 200mg/2mL in each single-dose PFS 200mg/2mL in each single-dose pen	Maintenance Dose: Inject 100mg subcutaneously at week 16 and every 8 weeks thereafter Inject 200mg subcutaneously at week 12 and every 4 weeks thereafter Inject 100mg subcutaneously every 8 weeks Inject 200mg subcutaneously every 4 weeks	1-month supply 3-month supply Other _____ Refills _____

Dispense needles, syringes, ancillary supplies and home medical equipment necessary to administer medication.

If shipped to physician's office, physician accepts on behalf of patient for administration in office.

Prescriber's signature required (sign below) (Physician attests this is his/her legal signature. NO STAMPS)**SIGN
HERE**

Date _____

Dispense as written _____

Date _____

Substitution allowed _____

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc.
Non-compliance with state-specific requirements could result in outreach to the prescriber.