

Please fax both pages of completed form to your team at 888.302.1028.

To reach your team, call toll-free 844.516.3319.

You can now monitor shipments and chat online if you have questions. Go to [MyAccredoPatients.com](http://MyAccredoPatients.com) to log in or get started.

Prescription & Enrollment Form

Ocrevus® (ocrelizumab)

accredo®

Four simple steps to submit your referral.

## 1 Patient Information



Please provide copies of front and back of all medical and prescription insurance cards.

New patient    Current patient

Patient's first name \_\_\_\_\_ Last name \_\_\_\_\_ Middle initial \_\_\_\_\_

Male    Female    Last 4 digits of SSN \_\_\_\_\_ Date of birth \_\_\_\_\_

Street address \_\_\_\_\_ Apt # \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Home phone \_\_\_\_\_ Cell phone \_\_\_\_\_ E-mail address \_\_\_\_\_

Parent/guardian (if applicable) \_\_\_\_\_

Home phone \_\_\_\_\_ Cell phone \_\_\_\_\_ E-mail address \_\_\_\_\_

Alternate caregiver/contact \_\_\_\_\_

Home phone \_\_\_\_\_ Cell phone \_\_\_\_\_ E-mail address \_\_\_\_\_

OK to leave message with alternate caregiver/contact

Patient's primary language:    English    Other    If other, please specify \_\_\_\_\_

## 2 Prescriber Information

All fields must be completed to expedite prescription fulfillment.

Date \_\_\_\_\_ Time \_\_\_\_\_ Date medication needed \_\_\_\_\_

Prescriber's first name \_\_\_\_\_ Last name \_\_\_\_\_

Prescriber's title \_\_\_\_\_ If NP or PA, under direction of Dr. \_\_\_\_\_

Office/Clinic/Institution name \_\_\_\_\_

Office address \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Phone \_\_\_\_\_ Fax \_\_\_\_\_ NPI # \_\_\_\_\_ License # \_\_\_\_\_

Office contact and title \_\_\_\_\_

Office contact phone number \_\_\_\_\_ Office contact e-mail \_\_\_\_\_

Clinic/institution name \_\_\_\_\_ Clinic/hospital affiliation \_\_\_\_\_

Street address \_\_\_\_\_ Suite # \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Infusion clinic contact name \_\_\_\_\_ E-mail address \_\_\_\_\_

Phone \_\_\_\_\_ Fax \_\_\_\_\_

**Note: Check the appropriate shipment options in Section 4: Prescribing Information.**

## 3 Clinical Information

Primary ICD-10 code:    Multiple Sclerosis: G35    Other \_\_\_\_\_    Laboratory results: LEVF \_\_\_\_\_

Platelets \_\_\_\_\_ Date \_\_\_\_\_ ANC \_\_\_\_\_ Date \_\_\_\_\_

Pregnancy test \_\_\_\_\_ (+/-) Date \_\_\_\_\_ Bilirubin \_\_\_\_\_ Date \_\_\_\_\_

FIRST TWO LOADING DOSES COMPLETED    Yes    No    Note: Ocrevus loading doses must be administered in a controlled setting.

EXPECTED DATE OF FIRST/NEXT INFUSION \_\_\_\_\_

NKDA    Known drug allergies \_\_\_\_\_

Concurrent meds \_\_\_\_\_

Patient's first name \_\_\_\_\_ Last name \_\_\_\_\_ Middle initial \_\_\_\_\_ Date of birth \_\_\_\_\_

Prescriber's first name \_\_\_\_\_ Last name \_\_\_\_\_ Phone \_\_\_\_\_

## 4 Prescribing Information

Medication	Dose	Directions	Quantity/Refills	Ship to*:
Ocrevus® (ocrelizumab) Initial dose (two infusions) <small>Note: Loading doses must be administered in a controlled infusion site.</small>	300mg/10mL SDV Vials are diluted in NS to a final concentration of 1.2mg/mL	<b>Infusion 1:</b> 300mg in 250mL of 0.9% NS. <b>Infusion 2 (2 weeks later):</b> 300mg in 250mL of 0.9% NS. Start infusion at 30mL per hour. Increase by 30mL per hour every 30 minutes. Maximum: 180mL per hour. Duration: 2.5 hours or longer.	Dispense: 2 vials No refills	Office Infusion Clinic Unknown
Ocrevus® (ocrelizumab) Subsequent doses (one infusion)	300mg/10mL SDV Vials are diluted in NS to a final concentration of 1.2mg/mL	Every 6 months infuse 600mg in 500mL of 0.9% NS. Start infusion at 40mL per hour. Increase by 40mL per hour every 30 minutes. Maximum: 200mL per hour. Duration: 3.5 hours or longer.	Dispense: 2 vials Refills 0 1	Home Office Infusion Clinic Unknown

All Ocrevus® orders to be administered via pump and peripheral line unless otherwise instructed.

### Additional Medication and Supplies for Home Infusion

#### Premedication Orders

Acetaminophen 650mg PO 30 min prior to infusion; Diphenhydramine 50mg PO 30 min prior to infusion; Methylprednisolone 100mg IV 30 min prior to infusion

Other \_\_\_\_\_

Send quantity sufficient for medication infusion  
All caregivers and ancillaries to be given per protocol from product package insert. (See next page).

#### Fluids for Reconstitution and Administration

0.9% NaCl 250mL x2 (initial dose); 0.9% NaCl 500mL (maintenance dose);  
0.9% NaCl Flush 10mL (3 mL pre- and post-infusion to maintain peripheral line patency)  
0.9% NACL 50mL  
0.9% NACL 100mL

If patient requires specific directions on additional medications or supplies, please provide change on the next page and sign.

#### Hypersensitivity/Anaphylaxis Orders\*

In the event of anaphylactic reaction, stop infusion of drug immediately. Start NS 15mL/hour; 0.9%NS 100mL.

Medicate with epinephrine pen auto-injector 0.3mg/0.3mL IM as needed for anaphylaxis. Call \*911\*, physician, or paramedic.

I authorize ancillary supplies or medical equipment necessary such as needles, syringes, etc. to administer the therapy as needed for administration.

Skilled nursing visit as needed to establish venous access, administer medication and assess general status and response to therapy.  
\*If nursing services will be required for therapy administration, the home health nurse will call for additional orders per state regulations.

If shipped to physician's office or infusion clinic, physician accepts on behalf of patient for administration in office or infusion clinic.

**Prescriber's signature (sign below) (Physician attests this is his/her legal signature. NO STAMPS)**

**SIGN  
HERE**

Date \_\_\_\_\_

Dispense as written \_\_\_\_\_

Date \_\_\_\_\_

Substitution allowed \_\_\_\_\_

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance with state-specific requirements could result in outreach to the prescriber.

Patient's first name \_\_\_\_\_ Last name \_\_\_\_\_ Middle initial \_\_\_\_\_ Date of birth \_\_\_\_\_

Prescriber's first name \_\_\_\_\_ Last name \_\_\_\_\_ Phone \_\_\_\_\_

## Accredo Additional Medications for Home Infusion Protocol as Per Package Insert

If your patient requires individualized dosing or administering, please cross out directions below, provide desired directions in the box and sign.

\_\_\_\_\_  
Date Signature

Medication	Dose	Directions
Diphenhydramine IV	50mg/1mL (25mg)	30 minutes prior to infusion, withdraw 0.5ml and inject into 50mL 0.9% NS. Infuse intravenously 101mL/hour over 30 min.
Diphenhydramine IV	50mg/1mL (50mg)	30 minutes prior to infusion, withdraw 1mL and inject into 50mL 0.9% NS. Infuse intravenously 102mL/hour over 30 min.
Methylprednisolone (Solu-Medrol) IV	100mg and Diphenhydramine PO	30 min prior to infusion, activate vial, withdraw 1.6mL/100mg, inject into 50mL 0.9% NS. Infuse intravenously 104mL/hour over 30 minutes.
Methylprednisolone (Solu-Medrol) IV	100mg and Diphenhydramine IV SIG	Activate vial, withdraw 1.6mL/100mg. Inject 100mg (1.6mL) intravenous push 0.2mL per minute for 8 minutes may increase to 0.4mL per minute for 4 minutes based on absence of infusion reactions (nausea, vomiting, headache, flushing, vital sign change) 30 minutes prior to Ocrevus.
Methylprednisolone (Solu-Medrol) IV	125mg SIG	30 minutes prior to infusion, activate vial, withdraw 2mL/125mg, inject into 100mL 0.9% NS. Infuse intravenously 204mL/hour over 30 minutes.
Methylprednisolone (Solu-Medrol) IV	250mg SIG	30 minutes prior to infusion, activate vial, withdraw 4mL/250mg, inject into 100mL 0.9% NS. Infuse intravenously 208mL/hour over 30 minutes.
Methylprednisolone (Solu-Medrol) IV	500mg SIG	30 min prior to infusion, activate vial, withdraw 8mL/500mg, inject into 100mL 0.9% NS. Infuse intravenously 216mL/hour over 30 minutes.
Methylprednisolone IV	125mg vial and Bacteriostatic water	Reconstitute Methylprednisolone 125mg with 2mL of Bacteriostatic water for injection. Withdraw 1.6mL/100mg. a. Inject 100mg (1.6mL) intravenous push 0.2mL per minute for 8 minutes may increase to 0.4mL per minute for 4 minutes based on absence of infusion reactions (nausea, vomiting, headache, flushing, vital sign change) 30 minutes prior to Ocrevus. b. Withdraw 1.6mL and inject into 50mL 0.9% NS. Infuse intravenously 104mL/hour over 30 minutes. 30 minutes prior to Ocrevus.
Famotidine IV	20mg	30 minutes prior to infusion, withdraw 2mL and inject into 100mL 0.9% NS. Infuse intravenously 204mL/hour over 30 minutes.
Famotidine IV	10mg	30 minutes prior to infusion, withdraw 1mL and inject into 100mL 0.9% NS. Infuse intravenously 202mL/hour over 30 minutes.