

Please fax both pages of completed form to your PAH team at 800.711.3526.

To reach your PAH team, call toll-free 888.200.2811, option 2, then option 1.

You can now monitor shipments and chat online if you have questions. Go to [MyAccredoPatients.com](http://MyAccredoPatients.com) to log in or get started.

Prescription & Enrollment Form  
**PAH Infusion**



Four simple steps to submit your referral.

# 1 Patient Information



Please attach copies of front and back of the patient's medical and prescription insurance cards.

New patient  Current patient

Patient's first name \_\_\_\_\_ Last name \_\_\_\_\_ Middle initial \_\_\_\_\_

Male  Female  Last 4 digits of SSN \_\_\_\_\_ Date of birth \_\_\_\_\_

Street address \_\_\_\_\_ Apt # \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Home phone \_\_\_\_\_ Cell phone \_\_\_\_\_ E-mail address \_\_\_\_\_

Parent/guardian (if applicable) \_\_\_\_\_

Home phone \_\_\_\_\_ Cell phone \_\_\_\_\_ E-mail address \_\_\_\_\_

Alternate caregiver/contact \_\_\_\_\_

Home phone \_\_\_\_\_ Cell phone \_\_\_\_\_ E-mail address \_\_\_\_\_

OK to leave message with alternate caregiver/contact

Patient's primary language: English  Other  If other, please specify \_\_\_\_\_

# 2 Prescriber Information

All fields must be completed to expedite prescription fulfillment.

Date \_\_\_\_\_ Time \_\_\_\_\_ Date medication needed \_\_\_\_\_

Prescriber's first name \_\_\_\_\_ Last name \_\_\_\_\_

Prescriber's title \_\_\_\_\_ If NP or PA, under direction of Dr. \_\_\_\_\_

Office address \_\_\_\_\_

Office contact and title \_\_\_\_\_

Office contact phone number \_\_\_\_\_ Office contact e-mail \_\_\_\_\_

Office/clinic/institution name \_\_\_\_\_ Clinic/hospital affiliation \_\_\_\_\_

Street address \_\_\_\_\_ Suite # \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Phone \_\_\_\_\_ Fax \_\_\_\_\_ NPI # \_\_\_\_\_ License # \_\_\_\_\_

Deliver product to: Office  Patient's home  Clinic  Clinic location \_\_\_\_\_

# 3 Clinical Information

Primary ICD-10 code (REQUIRED): \_\_\_\_\_

Diagnosis: ICD 127.0 - Pulmonary arterial hypertension (PAH) Idiopathic PAH Familial PAH  
ICD 127.21 - Pulmonary arterial hypertension Congenital heart disease

Connective tissue disease  HIV  Other \_\_\_\_\_

Concurrent meds \_\_\_\_\_

Weight \_\_\_\_\_ kg/lbs Height \_\_\_\_\_ cm/in Date recorded \_\_\_\_\_ Diabetic: Yes  No

NKDA  Known drug allergies \_\_\_\_\_

Select one: Urgent—Patient in hospital  Emergent—Admission within 48–72 hours  Standard—Admission after 4 days or more

Start-of-care date (REQUIRED) \_\_\_\_\_ Tentative discharge date \_\_\_\_\_

Discharge planner/coordinator name \_\_\_\_\_

Patient's first name \_\_\_\_\_ Last name \_\_\_\_\_ Middle initial \_\_\_\_\_ Date of birth \_\_\_\_\_

Prescriber's first name \_\_\_\_\_ Last name \_\_\_\_\_ Phone \_\_\_\_\_

## 4 Prescribing Information

The following prostacyclin therapies require additional information (e.g., diluent or titration). Please be sure to complete all information.

Medication	Pump and diluent	Dose and directions	Quantity/Refills
Flolan (epoprostenol)	CADD Legacy pH12 sterile diluent for Flolan	<b>Continuous IV infusion administered via ambulatory pump.</b> Initial dose _____ ng per kg per min. Dosing weight _____ kg. Titrate by _____ ng per kg per min every _____ days until _____ ng per kg per min is reached. Final concentration is _____ ng per mL. <hr/> <b>Continuous subcutaneous infusion administered via ambulatory pump.</b> Initial dose _____ ng per kg per min. Dosing weight _____ kg. Titrate by _____ ng per kg per min every _____ days until _____ ng per kg per min is reached. Final concentration is _____ ng per mL.	1-month supply 3-month supply Other _____ Refills _____
epoprostenol (generic Flolan)	CADD Legacy epoprostenol sterile diluent for injection		
epoprostenol (generic Veletri)	CADD Legacy 0.9% sodium chloride sterile water for injection		
treprostinil IV	CADD Legacy treprostinil sterile diluent for injection 0.9% sodium chloride sterile water for injection		
treprostinil IV	CADD MS3 0.9% sodium chloride sterile water for injection		
treprostinil subcut	CADD MS3		
Other instructions _____			
You must note the name of the brand product if brand is medically necessary for your patient _____			
Prescriber, please check here to authorize ancillary supplies such as needles, syringes, sterile water, infusion device, nebulizer, etc. to administer the therapy as needed for administration.			Send quantity sufficient for medication days supply
<b>Home nursing request to be provided by Accredo nursing staff (check all that apply)</b> In-hospital training (Accredo) Post-discharge visit/in-home follow-up Dispense teaching kits Home assessment/training prior to initiation of Flolan/treprostinil/Tyvaso therapy <b>DECLINE</b> all referenced nursing <i>If nursing services will be required for therapy administration, the home health nurse will call for additional orders per state regulations.</i>			

Prescriber's signature required (sign below) (Physician attests this is his/her legal signature. **NO STAMPS**)

**SIGN  
HERE**

Date \_\_\_\_\_

Dispense as written

Date \_\_\_\_\_

Substitution allowed

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance with state-specific requirements could result in outreach to the prescriber.