

Please fax all pages of completed form to your team at 833.951.1686.

To reach your team, call toll-free 800.442.5781.

You can now monitor shipments and chat online if you have questions. Go to [MyAccredoPatients.com](https://MyAccredoPatients.com) to log in or get started.

Prescription & Enrollment Form  
**Soliris® (eculizumab)**

accredo®

Four simple steps to submit your referral.

## 1 Patient Information



Please provide copies of front and back of all medical and prescription insurance cards.

New patient      Current patient

Patient's first name \_\_\_\_\_ Last name \_\_\_\_\_ Middle initial \_\_\_\_\_

Preferred patient first name \_\_\_\_\_ Preferred patient last name \_\_\_\_\_

Sex at birth:    Male    Female    Gender identity \_\_\_\_\_ Pronouns \_\_\_\_\_ Last 4 digits of SSN \_\_\_\_\_

Date of birth \_\_\_\_\_ Street address \_\_\_\_\_ Apt # \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Home phone \_\_\_\_\_ Cell phone \_\_\_\_\_ Email address \_\_\_\_\_

Parent/guardian (if applicable) \_\_\_\_\_

Home phone \_\_\_\_\_ Cell phone \_\_\_\_\_ Email address \_\_\_\_\_

Alternate caregiver/contact \_\_\_\_\_

Home phone \_\_\_\_\_ Cell phone \_\_\_\_\_ Email address \_\_\_\_\_

OK to leave message with alternate caregiver/contact

Patient's primary language:    English    Other    If other, please specify \_\_\_\_\_

## 2 Prescriber Information

All fields must be completed to expedite prescription fulfillment.

Date \_\_\_\_\_ Time \_\_\_\_\_ Date medication needed \_\_\_\_\_

Office/clinic/institution name \_\_\_\_\_

**Prescriber info:** Prescriber's first name \_\_\_\_\_ Last name \_\_\_\_\_

Prescriber's title \_\_\_\_\_ If NP or PA, under direction of Dr. \_\_\_\_\_

Office phone \_\_\_\_\_ Fax \_\_\_\_\_ NPI # \_\_\_\_\_ License # \_\_\_\_\_

Office contact and title \_\_\_\_\_ Office contact email \_\_\_\_\_

Office street address \_\_\_\_\_ Suite # \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Infusion location:    Patient's home    Prescriber's office    Infusion site    If infusion site, complete information below dotted line:  
-----

**Infusion info:** Infusion site name \_\_\_\_\_ Clinic/hospital affiliation \_\_\_\_\_

Site street address \_\_\_\_\_ Suite # \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Infusion site contact \_\_\_\_\_ Phone \_\_\_\_\_ Fax \_\_\_\_\_ Email \_\_\_\_\_

## 3 Clinical Information

**Primary ICD-10 code (REQUIRED):**    D59.5 Paroxysmal nocturnal hemoglobinuria    D59.3 Hemolytic-uremic syndrome    D59.32 Hereditary hemolytic-uremic syndrome    D59.39 Other hemolytic uremic syndrome    G70.00 Myasthenia Gravis without (acute) exacerbation    G70.01 Myasthenia Gravis with (acute) exacerbation    G36.0 Neuromyelitis optica    D58.8 Other specified hereditary hemolytic anemias    D59.4 Other non-autoimmune hemolytic anemias (including microangiopathic hemolytic anemia)

**D59.8 Other acquired hemolytic anemias**    Other \_\_\_\_\_

MG-ADL\* score (if known) \_\_\_\_\_ Weight \_\_\_\_\_ kg/lbs    Height \_\_\_\_\_ cm/in    Date recorded \_\_\_\_\_

NKDA    Known drug allergies \_\_\_\_\_

Concurrent meds \_\_\_\_\_

Adverse reactions with previous Soliris treatments? \_\_\_\_\_

Patient's first name \_\_\_\_\_ Last name \_\_\_\_\_ Middle initial \_\_\_\_\_ Date of birth \_\_\_\_\_

Prescriber's first name \_\_\_\_\_ Last name \_\_\_\_\_ Phone \_\_\_\_\_

## 4 Prescribing Information

Medication	Strength/Formulation	Directions
Soliris® (eculizumab)	300mg/30mL vial	<b>Loading dose:</b> _____ mg IV every _____ weeks for _____ weeks. <b>Maintenance dose:</b> _____ mg IV every _____ weeks. <b>Infusion method:</b> Gravity    Pump    Other _____
<b>Dilution and infusion rate</b>	<b>Loading dose:</b> Dilute Soliris with selected diluent as directed per manufacturer guidelines to a final concentration of 5mg/mL. If different, list here _____ <b>If adult patient:</b> Infusion rate: As directed per manufacturer guidelines _____ If different, list here _____ <b>If pediatric patient:</b> Infusion rate: As directed per manufacturer guidelines _____ If different, list here _____	<b>Maintenance dose:</b> Dilute Soliris with selected diluent as directed per manufacturer guidelines to a final concentration of 5mg/mL. If different, list here _____ <b>If adult patient:</b> Infusion rate: As directed per manufacturer guidelines _____ If different, list here _____ <b>If pediatric patient:</b> Infusion rate: As directed per manufacturer guidelines _____ If different, list here _____
<b>Check one (0.9% Sodium Chloride will be used if no box is checked):</b> 0.9% Sodium Chloride Injection    0.45% Sodium Chloride Injection    5% Dextrose in Water Injection    Ringer's Injection		
Other instructions _____		
Has the patient received Meningitis vaccination?    Yes    No <b>MenACWY</b> <b>1st Dose</b> Brand:    Menveo    Menactra    MenQuadFi    Other/Unknown Date of administration: _____ <b>2nd Dose</b> Brand:    Menveo    Menactra    MenQuadFi    Other/Unknown Date of administration: _____ <b>MenABCWY</b> <b>1st Dose</b> Brand:    Penbraya    Other/Unknown Date of administration: _____ <b>2nd Dose</b> Brand:    Penbraya    Other/Unknown Date of administration: _____		
<b>MenB</b> <b>1st Dose</b> Brand:    Bexsero    Trumenba    Other/Unknown Date of administration: _____ <b>2nd Dose</b> Brand:    Bexsero    Trumenba    Other/Unknown Date of administration: _____ <b>3rd Dose</b> Brand:    Bexsero    Trumenba    Other/Unknown Date of administration: _____ <b>Antibacterial Drug Prophylaxis (if applicable)</b> Did patient receive antibacterial drug prophylaxis?    Yes    No Start Date: _____		
<b>Complete the below section if assistance from Accredo is requested in the coordination of your patient's infusion therapy</b>		
Is Accredo home nursing service requested?    Yes    No    Vascular access:    Peripheral    Central    Port		
Supplies: (please strike through if not required) Dispense needles, syringes, ancillary supplies and home medical equipment necessary to administer medication. <b>PERIPHERAL Access:</b> 0.9% Normal Saline 3mL intravenous before and after infusion, or as needed for line patency. If different, please list here _____ <b>PORT/CENTRAL Access:</b> 0.9% Normal Saline 5mL intravenous before and after infusion, or as needed for line patency. Heparin 10 units per mL 5mL intravenous as needed for final flush. If different, please list here _____		
Is your patient new to therapy?    Yes    No		

This form is a generic referral form that could be utilized for any Soliris provider and is meant to provide the pertinent information needed to process a Soliris referral.

If nursing services will be required for the therapy administration, the home health nurse will call for additional orders per state regulations.

ALL fields must be completed to expedite prescription fulfillment.

If shipped to physician's office or infusion clinic, physician accepts on behalf of patient for administration in office or infusion clinic.

**Prescriber's signature required (sign below) (Physician attests this is his/her legal signature. NO STAMPS)**SIGN  
HERE

Date

Dispense as written

Date

Substitution allowed

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc.  
 Non-compliance with state-specific requirements could result in outreach to the prescriber.

Patient's first name \_\_\_\_\_ Last name \_\_\_\_\_ Middle initial \_\_\_\_\_ Date of birth \_\_\_\_\_

Prescriber's first name \_\_\_\_\_ Last name \_\_\_\_\_ Phone \_\_\_\_\_

4 Prescribing Information

**Hypersensitivity/Anaphylaxis** Stop infusion  
**Medicate with:** Epinephrine 0.3mg Auto Injector – Stop infusion and inject dose per packaging for hypersensitivity/anaphylaxis (patient weighs greater than or equal to 30kg) **OR** Epinephrine JR 0.15mg/0.3mL Auto injector – Stop infusion and inject dose per packaging for hypersensitivity/anaphylaxis (patient weighs 15kg to 29kg)

**Premedications** *(Prescriber, please list any premedication(s) you want your patient to have.)*  
Drug \_\_\_\_\_ Directions \_\_\_\_\_  
Drug \_\_\_\_\_ Directions \_\_\_\_\_

**Quantity/Refills:** Dispense quantity sufficient for medication days supply for loading dose, then 1 month ongoing for maintenance dose. Refill x 1 year.  
Other \_\_\_\_\_

Skilled nursing visit as needed to establish venous access, administer medication and assess general status and response to therapy. If nursing services will be required for therapy administration, the home health nurse will call for additional orders per state regulations.

This form is a generic referral form that could be utilized for any Soliris provider and is meant to provide the pertinent information needed to process a Soliris referral. If nursing services will be required for the therapy administration, the home health nurse will call for additional orders per state regulations. ALL fields must be completed to expedite prescription fulfillment. If shipped to physician's office or infusion clinic, physician accepts on behalf of patient for administration in office or infusion clinic.

Prescriber's signature required (sign below) (Physician attests this is his/her legal signature. NO STAMPS)

SIGN HERE

Date

Dispense as written

Date

Substitution allowed

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance with state-specific requirements could result in outreach to the prescriber.