

Please fax both pages of completed form to your team at 888.302.1028.

To reach your team, call toll-free 844.516.3319.

You can now monitor shipments and chat online if you have questions. Go to [MyAccredoPatients.com](http://MyAccredoPatients.com) to log in or get started.

Prescription & Enrollment Form

**Spevigo<sup>®</sup> (spesolimab-sbzo)**

*accredo<sup>®</sup>*

Four simple steps to submit your referral.

## 1 Patient Information



Please provide copies of front and back of all medical and prescription insurance cards.

New patient    Current patient

Patient's first name \_\_\_\_\_ Last name \_\_\_\_\_ Middle initial \_\_\_\_\_

Sex at birth: Male Female Preferred pronouns \_\_\_\_\_ Last 4 digits of SSN \_\_\_\_\_ Date of birth \_\_\_\_\_

Street address \_\_\_\_\_ Apt # \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Home phone \_\_\_\_\_ Cell phone \_\_\_\_\_ E-mail address \_\_\_\_\_

Parent/guardian (if applicable) \_\_\_\_\_

Home phone \_\_\_\_\_ Cell phone \_\_\_\_\_ E-mail address \_\_\_\_\_

Alternate caregiver/contact \_\_\_\_\_

Home phone \_\_\_\_\_ Cell phone \_\_\_\_\_ E-mail address \_\_\_\_\_

OK to leave message with alternate caregiver/contact

Patient's primary language: English Other If other, please specify \_\_\_\_\_

## 2 Prescriber Information

All fields must be completed to expedite prescription fulfillment.

Date \_\_\_\_\_ Time \_\_\_\_\_ Date medication needed \_\_\_\_\_

Office/clinic/institution name \_\_\_\_\_

**Prescriber info:** Prescriber's first name \_\_\_\_\_ Last name \_\_\_\_\_

Prescriber's title \_\_\_\_\_ If NP or PA, under direction of Dr. \_\_\_\_\_

Office phone \_\_\_\_\_ Fax \_\_\_\_\_ NPI # \_\_\_\_\_ License # \_\_\_\_\_

Office contact and title \_\_\_\_\_ Office contact e-mail \_\_\_\_\_

Office street address \_\_\_\_\_ Suite # \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Infusion location: Patient's home Prescriber's office Infusion site If infusion site, complete information below dotted line: \_\_\_\_\_

**Infusion info:** Infusion site name \_\_\_\_\_ Clinic/hospital affiliation \_\_\_\_\_

Site street address \_\_\_\_\_ Suite # \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Infusion site contact \_\_\_\_\_ Phone \_\_\_\_\_ Fax \_\_\_\_\_ E-mail \_\_\_\_\_

## 3 Clinical Information

**Primary ICD-10 code (REQUIRED):** \_\_\_\_\_

NKDA Known drug allergies \_\_\_\_\_

Concurrent meds \_\_\_\_\_

Patient's first name \_\_\_\_\_ Last name \_\_\_\_\_ Middle initial \_\_\_\_\_ Date of birth \_\_\_\_\_

Prescriber's first name \_\_\_\_\_ Last name \_\_\_\_\_ Phone \_\_\_\_\_

## 4 Prescribing Information

**INFUSION LOCATION:** Patient's home    Healthcare facility

Medication	Strength/Formulation	Directions	Quantity/Refills
Spevigo® (spesolimab-sbzo)	450mg/7.5mL vial	Infuse 900mg (Two 450mg single dose vials) Intravenously once over 90 minutes	2 vials Refills _____

**Required medication and supplies for home infusion (please complete this section for home infusions only)**

<p><b>Premedication Orders</b>                  Acetaminophen 650mg PO 30 min prior to infusion;    Diphenhydramine 50mg PO 30 min prior to infusion                  Other _____</p>	Send quantity and refills sufficient for medication days supply
<p><b>Infusion method:</b> Gravity (Pediatric patients will be given a pump unless noted otherwise)</p>	
<p><b>Fluids for administration and reconstitution (please strike through if not required)</b>                  NS 0.9% 100mL                  NS 0.9% Flush (if central venous access, sterile flush will be provided)                  Choose administration access:    Peripheral access    Central venous access                  If central venous access: Flush with 10mL Sterile NS 0.9% before and after infusion.    Follow with heparin 100units/mL 5mL final flush                  If peripheral access: Flush with 3mL NS 0.9% before and after infusion and as needed</p>	
<p><b>Hypersensitivity/Anaphylaxis</b>                  Stop infusion  <b>Medicate with:</b>                  Epinephrine/EpiPen 0.3mg IM as needed for anaphylaxis (for children less than 30kg: Epinephrine 0.15mg)                  Start NS 0.9% 100mL at TKO    Diphenhydramine 50mg slow IVP PRN anaphylaxis                  Hydrocortisone 100mg slow IVP PRN anaphylaxis                  Methylprednisolone 125mg slow IVP PRN anaphylaxis    Diphenhydramine 50mg PO PRN anaphylaxis                  Other _____</p>	
Skilled nursing visit as needed to establish venous access, administer medication and assess general status and response to therapy. *If nursing services will be required for therapy administration, the home health nurse will call for additional orders per state regulations. Lab orders _____ Frequency _____	

Dispense needles, syringes, ancillary supplies and home medical equipment necessary to administer medication.

If shipped to physician's office, physician accepts on behalf of patient for administration in office.

**Prescriber's signature required (sign below) (Physician attests this is his/her legal signature. NO STAMPS)**

**SIGN  
HERE**

Date

Dispense as written

Date

Substitution allowed

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance with state-specific requirements could result in outreach to the prescriber.