

Please fax both pages of completed form to your team at 888.302.1028.

To reach your team, call toll-free 844.516.3319.

You can now monitor shipments and chat online if you have questions. Go to MyAccredoPatients.com to log in or get started.

Prescription & Enrollment Form Uplizna[®] (inebilizumab injection)

accredo[®]

Four simple steps to submit your referral.

1 Patient Information



Please provide copies of front and back of all medical and prescription insurance cards.

New patient Current patient

Patient's first name _____ Last name _____ Middle initial _____

Sex at birth: Male Female Preferred pronouns _____ Last 4 digits of SSN _____ Date of birth _____

Street address _____ Apt # _____

City _____ State _____ Zip _____

Home phone _____ Cell phone _____ E-mail address _____

Parent/guardian (if applicable) _____

Home phone _____ Cell phone _____ E-mail address _____

Alternate caregiver/contact _____

Home phone _____ Cell phone _____ E-mail address _____

OK to leave message with alternate caregiver/contact

Patient's primary language: English Other If other, please specify _____

2 Prescriber Information

All fields must be completed to expedite prescription fulfillment.

Date _____ Time _____ Date medication needed _____

Office/clinic/institution name _____

Prescriber info: Prescriber's first name _____ Last name _____

Prescriber's title _____ If NP or PA, under direction of Dr. _____

Office phone _____ Fax _____ NPI # _____ License # _____

Office contact and title _____ Office contact e-mail _____

Office street address _____ Suite # _____

City _____ State _____ Zip _____

Infusion location: Patient's home Prescriber's office Infusion site If infusion site, complete information below dotted line:

Infusion info: Infusion site name _____ Clinic/hospital affiliation _____

Site street address _____ Suite # _____

City _____ State _____ Zip _____

Infusion site contact _____ Phone _____ Fax _____ E-mail _____

3 Clinical Information

Primary ICD-10 code (REQUIRED): _____ **Diagnosis** G36.0 Neuromyelitis optica Other _____

Is the patient anti-aquaporin-4 antibody positive? Yes No Test pending

Prior NSMOD therapies tried/failed _____

Hep B vaccination: Yes No Date _____ Does the patient have active Hepatitis B infection? Yes No

Hepatitis B screening: Hepatitis B surface antigen (HBsAg) results Positive Negative Date _____

HB core antibody [HBcAb+] results Positive Negative Date _____

Does the patient have active or latent TB infection? Yes No Tuberculosis screening: Positive Negative Date _____

FIRST TWO LOADING DOSES COMPLETED Yes No **Note: Uplizna loading doses must be administered in a controlled setting.**

EXPECTED DATE OF FIRST/NEXT INFUSION _____

NKDA Known drug allergies _____

Concurrent meds _____

Patient's first name _____ Last name _____ Middle initial _____ Date of birth _____

Prescriber's first name _____ Last name _____ Phone _____

4 Prescribing Information

***Provide address for the selected shipment option.
Check Unknown if assistance is needed to identify infusion site.**

Medication	Dose	Directions	Quantity/Refills	Ship to*:
Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a controlled infusion site.	100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL	Infusion 1: 300mg in 250mL of 0.9% NS. Infusion 2 (2 weeks later): 300mg in 250mL of 0.9% NS. Start infusion at 42mL per hour for the first 30 minutes, increase to 125mL per hour for the next 30 minutes, then increase to 333mL per hour until finished. Duration: 90 minutes or longer Monitor patient for at least one hour after infusion completion for infusion reaction.	Dispense: 6 vials No refills	Office Infusion Clinic Unknown
Uplizna® (inebilizumab injection) Maintenance dose (one infusion)	100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL	Every 6 months (from first infusion) infuse 300mg in 250mL of 0.9% NS. Start infusion at 42mL per hour for the first 30 minutes, increase to 125mL per hour for the next 30 minutes, then increase to 333mL per hour until finished. Duration: 90 minutes or longer Monitor patient for at least one hour after infusion completion for infusion reaction.	Dispense: 3 vials Refills 0 1	Home Office Infusion Clinic Unknown

All Uplizna® orders to be administered via pump and peripheral line unless otherwise instructed.

Additional Medication and Supplies for Home Infusion

Premedication Orders

Acetaminophen 650mg PO 30 min prior to infusion; Diphenhydramine 50mg PO 30 min prior to infusion; Methylprednisolone 100mg IV 30 min prior to infusion

Other _____

Send quantity sufficient for medication infusion
All caregivers and ancillaries to be given per protocol from product package insert. (See next page).

Fluids for Reconstitution and Administration

0.9% NaCl 250mL x2 (initial dose); 0.9% NaCl 250mL (maintenance dose);
0.9% NaCl Flush 10mL (3 mL pre- and post-infusion to maintain peripheral line patency)
0.9% NACL 50mL
0.9% NACL 100mL

If patient requires specific directions on additional medications or supplies, please provide change on the next page and sign.

Hypersensitivity/Anaphylaxis Orders*

In the event of anaphylactic reaction, stop infusion of drug immediately. Start NS 15mL/hour; 0.9%NS 100mL. Medicate with epinephrine pen auto-injector 0.3mg/0.3mL IM as needed for anaphylaxis. Call *911*, physician, or paramedic.

I authorize ancillary supplies or medical equipment necessary such as needles, syringes, etc. to administer the therapy as needed for administration.

Skilled nursing visit as needed to establish venous access, administer medication and assess general status and response to therapy.
*If nursing services will be required for therapy administration, the home health nurse will call for additional orders per state regulations.

If shipped to physician's office or infusion clinic, physician accepts on behalf of patient for administration in office or infusion clinic.

Prescriber's signature required (sign below) (Physician attests this is his/her legal signature. NO STAMPS)

**SIGN
HERE**

Date

Dispense as written

Date

Substitution allowed

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance with state-specific requirements could result in outreach to the prescriber.



The document(s) accompanying this transmission may contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender immediately and arrange for the return or destruction of these documents. All rights in the product names, trade names or logos of all third-party products that appear in this form, whether or not appearing with the trademark symbol, belong exclusively to their respective owners. © 2022 Accredo Health Group, Inc. I An Express Scripts Company. All rights reserved. MSS-00009-112222 CRP2310_1765

Accredo Additional Medications for Home Infusion Protocol as Per Package Insert

If your patient requires individualized dosing or administering, please cross out directions below, provide desired directions in the box and sign.

Date	Signature																																		
		<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 25%;">Medication</th> <th style="width: 25%;">Dose</th> <th style="width: 50%;">Directions</th> </tr> </thead> <tbody> <tr> <td>Diphenhydramine</td> <td>50mg/1mL (25mg)</td> <td>30 minutes prior to infusion, withdraw 0.5ml and inject into 50mL 0.9% NS. Infuse intravenously 101mL/hour over 30 min.</td> </tr> <tr> <td>Diphenhydramine</td> <td>50mg/1mL (50mg)</td> <td>30 minutes prior to infusion, withdraw 1mL and inject into 50mL 0.9% NS. Infuse intravenously 102mL/hour over 30 min.</td> </tr> <tr> <td>Methylprednisolone</td> <td>100mg and Benadryl PO</td> <td>30 min prior to infusion, activate vial, withdraw 1.6mL/100mg, inject into 50mL 0.9% NS. Infuse intravenously 104mL/hour over 30 minutes.</td> </tr> <tr> <td>Methylprednisolone</td> <td>100mg and Benadryl IV SIG</td> <td>Activate vial, withdraw 1.6mL/100mg. Inject 100mg (1.6mL) intravenous push 0.2mL per minute for 8 minutes may increase to 0.4mL per minute for 4 minutes based on absence of infusion reactions (nausea, vomiting, headache, flushing, vital sign change) 30 minutes prior to Uplizna.</td> </tr> <tr> <td>Methylprednisolone</td> <td>125mg SIG</td> <td>30 minutes prior to infusion, activate vial, withdraw 2mL/125mg, inject into 100mL 0.9% NS. Infuse intravenously 204mL/hour over 30 minutes.</td> </tr> <tr> <td>Methylprednisolone</td> <td>250mg SIG</td> <td>30 minutes prior to infusion, activate vial, withdraw 4mL/250mg, inject into 100mL 0.9% NS. Infuse intravenously 208mL/hour over 30 minutes.</td> </tr> <tr> <td>Methylprednisolone</td> <td>500mg SIG</td> <td>30 min prior to infusion, activate vial, withdraw 8mL/500mg, inject into 100mL 0.9% NS. Infuse intravenously 216mL/hour over 30 minutes.</td> </tr> <tr> <td>Methylprednisolone</td> <td>125mg vial and Bacteriostatic water</td> <td> Reconstitute Methylprednisolone 125mg with 2mL of Bacteriostatic water for injection. Withdraw 1.6mL/100mg. <ol style="list-style-type: none"> a. Inject 100mg (1.6mL) intravenous push 0.2mL per minute for 8 minutes may increase to 0.4mL per minute for 4 minutes based on absence of infusion reactions (nausea, vomiting, headache, flushing, vital sign change) 30 minutes prior to Uplizna. b. Withdraw 1.6mL and inject into 50mL 0.9% NS. Infuse intravenously 104mL/hour over 30 minutes. 30 minutes prior to Uplizna. </td> </tr> <tr> <td>Famotidine</td> <td>20mg</td> <td>30 minutes prior to infusion, withdraw 2mL and inject into 100mL 0.9% NS. Infuse intravenously 204mL/hour over 30 minutes.</td> </tr> <tr> <td>Famotidine</td> <td>10mg</td> <td>30 minutes prior to infusion, withdraw 1mL and inject into 100mL 0.9% NS. Infuse intravenously 202mL/hour over 30 minutes.</td> </tr> </tbody> </table>	Medication	Dose	Directions	Diphenhydramine	50mg/1mL (25mg)	30 minutes prior to infusion, withdraw 0.5ml and inject into 50mL 0.9% NS. Infuse intravenously 101mL/hour over 30 min.	Diphenhydramine	50mg/1mL (50mg)	30 minutes prior to infusion, withdraw 1mL and inject into 50mL 0.9% NS. Infuse intravenously 102mL/hour over 30 min.	Methylprednisolone	100mg and Benadryl PO	30 min prior to infusion, activate vial, withdraw 1.6mL/100mg, inject into 50mL 0.9% NS. Infuse intravenously 104mL/hour over 30 minutes.	Methylprednisolone	100mg and Benadryl IV SIG	Activate vial, withdraw 1.6mL/100mg. Inject 100mg (1.6mL) intravenous push 0.2mL per minute for 8 minutes may increase to 0.4mL per minute for 4 minutes based on absence of infusion reactions (nausea, vomiting, headache, flushing, vital sign change) 30 minutes prior to Uplizna.	Methylprednisolone	125mg SIG	30 minutes prior to infusion, activate vial, withdraw 2mL/125mg, inject into 100mL 0.9% NS. Infuse intravenously 204mL/hour over 30 minutes.	Methylprednisolone	250mg SIG	30 minutes prior to infusion, activate vial, withdraw 4mL/250mg, inject into 100mL 0.9% NS. Infuse intravenously 208mL/hour over 30 minutes.	Methylprednisolone	500mg SIG	30 min prior to infusion, activate vial, withdraw 8mL/500mg, inject into 100mL 0.9% NS. Infuse intravenously 216mL/hour over 30 minutes.	Methylprednisolone	125mg vial and Bacteriostatic water	Reconstitute Methylprednisolone 125mg with 2mL of Bacteriostatic water for injection. Withdraw 1.6mL/100mg. <ol style="list-style-type: none"> a. Inject 100mg (1.6mL) intravenous push 0.2mL per minute for 8 minutes may increase to 0.4mL per minute for 4 minutes based on absence of infusion reactions (nausea, vomiting, headache, flushing, vital sign change) 30 minutes prior to Uplizna. b. Withdraw 1.6mL and inject into 50mL 0.9% NS. Infuse intravenously 104mL/hour over 30 minutes. 30 minutes prior to Uplizna. 	Famotidine	20mg	30 minutes prior to infusion, withdraw 2mL and inject into 100mL 0.9% NS. Infuse intravenously 204mL/hour over 30 minutes.	Famotidine	10mg	30 minutes prior to infusion, withdraw 1mL and inject into 100mL 0.9% NS. Infuse intravenously 202mL/hour over 30 minutes.
Medication	Dose	Directions																																	
Diphenhydramine	50mg/1mL (25mg)	30 minutes prior to infusion, withdraw 0.5ml and inject into 50mL 0.9% NS. Infuse intravenously 101mL/hour over 30 min.																																	
Diphenhydramine	50mg/1mL (50mg)	30 minutes prior to infusion, withdraw 1mL and inject into 50mL 0.9% NS. Infuse intravenously 102mL/hour over 30 min.																																	
Methylprednisolone	100mg and Benadryl PO	30 min prior to infusion, activate vial, withdraw 1.6mL/100mg, inject into 50mL 0.9% NS. Infuse intravenously 104mL/hour over 30 minutes.																																	
Methylprednisolone	100mg and Benadryl IV SIG	Activate vial, withdraw 1.6mL/100mg. Inject 100mg (1.6mL) intravenous push 0.2mL per minute for 8 minutes may increase to 0.4mL per minute for 4 minutes based on absence of infusion reactions (nausea, vomiting, headache, flushing, vital sign change) 30 minutes prior to Uplizna.																																	
Methylprednisolone	125mg SIG	30 minutes prior to infusion, activate vial, withdraw 2mL/125mg, inject into 100mL 0.9% NS. Infuse intravenously 204mL/hour over 30 minutes.																																	
Methylprednisolone	250mg SIG	30 minutes prior to infusion, activate vial, withdraw 4mL/250mg, inject into 100mL 0.9% NS. Infuse intravenously 208mL/hour over 30 minutes.																																	
Methylprednisolone	500mg SIG	30 min prior to infusion, activate vial, withdraw 8mL/500mg, inject into 100mL 0.9% NS. Infuse intravenously 216mL/hour over 30 minutes.																																	
Methylprednisolone	125mg vial and Bacteriostatic water	Reconstitute Methylprednisolone 125mg with 2mL of Bacteriostatic water for injection. Withdraw 1.6mL/100mg. <ol style="list-style-type: none"> a. Inject 100mg (1.6mL) intravenous push 0.2mL per minute for 8 minutes may increase to 0.4mL per minute for 4 minutes based on absence of infusion reactions (nausea, vomiting, headache, flushing, vital sign change) 30 minutes prior to Uplizna. b. Withdraw 1.6mL and inject into 50mL 0.9% NS. Infuse intravenously 104mL/hour over 30 minutes. 30 minutes prior to Uplizna. 																																	
Famotidine	20mg	30 minutes prior to infusion, withdraw 2mL and inject into 100mL 0.9% NS. Infuse intravenously 204mL/hour over 30 minutes.																																	
Famotidine	10mg	30 minutes prior to infusion, withdraw 1mL and inject into 100mL 0.9% NS. Infuse intravenously 202mL/hour over 30 minutes.																																	